

You should read this labeling carefully before use.
Please keep it so that you can read it when you need to.

Schedule II
pharmaceuticals



Therapeutic Agent for Eczema/Dermatitis

プレバリン[®]α クイック 軟膏

Prevaline α Quick Ointment

Effective in the affected area, low activity in the body

[Characteristics]

Prevaline α Quick Ointment is an antedrug type steroid that shows excellent effects on eczema/dermatitis.

Skin diseases, such as eczema, dermatitis, and heat rash, are common, but if itching or rash is strong, the skin may be scratched unconsciously, which further worsens the symptoms.

Prevaline α Quick Ointment is an ointment-type drug for treating skin diseases to protect the affected area by combining a steroid to suppress such inflammation and an ingredient to suppress itching.

An antedrug steroid is:

The steroid component contained in this product, prednisolone valerate acetate, is an antedrug steroid that works effectively on the skin surface, is rapidly broken down while being absorbed into the body, and becomes an ingredient with low activity.



Precautions

When not to use the product



(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

1. **The medicine should not be used on the following parts:**
 - (1) Varicella (chickenpox), athlete's foot/tinea or purulent lesions
 - (2) Periorbital area and mucosa (e.g., lips)
2. **Do not apply to the face extensively.**
3. **Do not use this medicine for a long time.**



Consultation

1. **The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use:**
 - (1) Patients undergoing medical treatment from a physician
 - (2) Pregnant women or women suspected of being pregnant
 - (3) Patients who have experienced allergic symptoms associated with drugs, etc.
 - (4) Those with extensive lesions
 - (5) Those with severe moistening or erosion
2. **If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use, and show this document to your physician, pharmacist, or registered salesperson for a consultation.**

Related sites	Symptoms
Skin	Rash/redness, itching, swelling
Skin (affected area)	Tinea, such as athlete's foot/ringworm, acne, purulent symptom, persistent irritation

3. **If symptoms do not improve even after using this medicine for 5 to 6 days, stop using this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.**

(Be sure to read the reverse side)

[Indications]

Eczema, dermatitis, heat rash, contact dermatitis, itching, insect bites, urticaria




[Dosage and administration]

Apply an appropriate amount to the affected lesion several times a day.

< Precautions of Dosage and Administration >

- (1) When allowing children to use the medicine, a guardian must be present to watch and tell them how to use it.
- (2) Attention should be paid so that the medicine will not come into contact with eyes. If the medicine comes into contact with eyes, immediately rinse them with water or lukewarm water. If the symptoms are severe, seek an ophthalmologist for medical treatment.
- (3) This medication is only for external use.
- (4) After applying this product, do not cover the affected area with unbreathable materials such as plastic wrap film.

[How to make a hole in the tube]



Turn the cap upside down, and press the protrusion firmly against the tip of the tube to release it.

[Ingredients] Per 100 g

Prednisolone valerate acetate	0.15g
Chlorpheniramine maleate	1 g
Vitamin E acetate	0.5 g
Lidocaine	2 g
Isopropyl methylphenol	0.1g
Allantoin	0.25g
γ-oryzanol	1 g

Inactive ingredients: Stearyl alcohol, cetanol, liquid paraffin, white petrolatum, medium-chain triglyceride, microcrystalline wax, squalane, diethyl sebacate, isopropyl palmitate

[Precautions for storage and handling]

- (1) Store in a cool, low humidity place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)
- (4) Do not use the product after the expiration date.

Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.