

Product name: **ヘパリーゼ®ドリンクII** **Hepalyse Drink II**



[Precautions] **Consultation**

Class 3 drug

1. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Related site	Symptoms
Skin	Rash
Gastrointestinal system	Heartburn, nausea, diarrhea

2. When symptoms do not improve even after taking the medicine for a while, stop taking the preparation and consult a physician, pharmacist, or registered salesperson, being sure to take this instruction leaflet with you.

[Indications] Nutritional fortification, nutritional supplement in cases of gastrointestinal disorders, nutritional disorders, during and after illness, physical fatigue, febrile debilitating diseases, and during pregnancy and lactation, and weak constitution

[Dosage and administration] Adults (15 years or older): Take 1 bottle (50 mL) once daily. Persons younger than 15 years must not take this drug.

- Follow the dosage and administration.

[Ingredients] Content of 1 bottle (50 mL): 200 mg of liver hydrolysate, 120 mg of sodium chondroitin sulfate, 20 mg of diisopropylamine dichloroacetate, 54 mg of schisandra fruit extract (equivalent to 300 mg of schisandra fruit), 0.2 mL of lycium fruit fluid extract (equivalent to 200 mg of lycium fruit), 0.1 mL of astragalus root fluid extract (equivalent to 100 mg of astragalus root), 1000 mg of taurine, 5 mg of vitamin B₂ phosphate ester, 10 mg of vitamin B₆, and 20 mg of nicotinamide

Excipients: Isomerized sugar, caramel, sodium benzoate, butylparaben, sodium citrate hydrate, fragrance, ethanol, propylene glycol, vanillin, ethylvanillin, 2 pH modifiers (alcohol ≤ 0.1 mL)

- Urine may turn yellow because of the Vitamin B₂ phosphate contained in this drug.

[Precautions for storage and handling] (1) Store in a cool place away from direct sunlight. (2) Keep out of reach of children. (3) On rare occasions, this drug may become turbid; however, its efficacy is unchanged when this occurs. (4) Do not take this product after the expiration date.

G16

Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.